

WARNING:
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

CATOLYST®

FOR THE TREATMENT AND PREVENTION OF COCCIDIOSIS IN CATTLE UP
TO 9 MONTHS CAUSED BY *EIMERIA BOVIS* OR *EIMERIA ZUERNII*.

Active ingredient: 50 g/L Toltrazuril



NET CONTENT: 2.5L

PLCAT2.5F-01

CATOLYST®

CATOLYST® controls coccidiosis caused by *Eimeria bovis* or *Eimeria zuernii* in young cattle. Coccidiosis typically occurs in cattle less than a year old causing diarrhoea which is often blood stained. Recently weaned calves are particularly susceptible to outbreaks of clinical disease. CATOLYST® is effective in one dose because it attacks all stages of the parasite in the animal.

DIRECTIONS FOR USE By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label. Thoroughly shake container before use.

CATTLE DOSE: Give 3 mls of CATOLYST® per 10kg of bodyweight by mouth. For the treatment of clinical disease treat all affected and in contact animals.

For prevention of coccidiosis on farms with a known history treat prior to the expected onset of clinical signs. For prevention of coccidiosis in recently weaned calves treat at weaning time when meal feeding ceases.

GENERAL INSTRUCTIONS: Before using this product obtain a diagnosis for diarrhoea. Early treatment of calves as soon as clinical signs of disease are seen and confirmed to be due to coccidiosis will reduce the impact of an outbreak by preventing further intestinal damage in affected animals and preventing non-affected animals from developing diarrhoea. To obtain maximum benefit on farms with a history of coccidiosis, CATOLYST® needs to be given approximately 1 week prior to the expected onset of clinical signs. Treatment of newly weaned calves at the time of meal withdrawal will control coccidiosis associated with weaning.

To achieve effective control of coccidiosis good management and hygiene are vital. In animals already showing signs of diarrhoea, additional supportive therapy may be required. Where necessary seek advice from a veterinarian or animal health adviser regarding effective management and control of coccidiosis.

WARNING: Suspected of damaging fertility or the unborn child. May cause damage to organs. **Read label before use.** Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Use personal protective equipment as required. Do not eat, drink or smoke when using this product. May cause drowsiness or dizziness. May cause respiratory irritation. Do not breathe vapours. Use only outdoors or in a well-ventilated area. Wash hands thoroughly after handling. **FIRST AID:** Call a POISON CENTER (0800 764 766) or doctor/physician if you feel unwell. If exposed or concerned get medical advice/attention. **IF INHALED:** Remove to fresh air and keep at rest in a position comfortable for breathing. **STORAGE:** Store locked up. Store in a well-ventilated place. Keep container tightly closed. **DISPOSAL:** Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility.

WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.


Cattle producing meat and offal for human consumption must not be sold for slaughter during or within 56 days of the last treatment.

ACVM Registration No. A011436
See www.foodsafety.govt.nz for registration conditions

PLCAT2.5B-01
STORE below 25°C
B.
EXP.



Registered to
Alleva Animal Health Ltd
15 Calman Place,
Birkenhead, Auckland



CATOLYST® is a registered trademark of Alleva Animal Health

WARNING:
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

CATOLYST®

FOR THE TREATMENT AND PREVENTION OF
COCCIDIOSIS IN CATTLE UP TO 9 MONTHS CAUSED
BY *EIMERIA BOVIS* OR *EIMERIA ZUERNII*.

Active ingredient: 50 g/L Toltrazuril



NET CONTENT: 1L

PLCAT1F-01

GENERAL INSTRUCTIONS

Before using this product obtain a diagnosis for diarrhoea. Early treatment of calves as soon as clinical signs of disease are seen and confirmed to be due to coccidiosis will reduce the impact of an outbreak by preventing further intestinal damage in affected animals and preventing non-affected animals from developing diarrhoea.

To obtain maximum benefit on farms with a history of coccidiosis, CATOLYST® needs to be given approximately 1 week prior to the expected onset of clinical signs. Treatment of newly weaned calves at the time of meal withdrawal will control coccidiosis associated with weaning.

To achieve effective control of coccidiosis good management and hygiene are vital. In animals already showing signs of diarrhoea, additional supportive therapy may be required. Where necessary seek advice from a veterinarian or animal health adviser regarding effective management and control of coccidiosis.

WARNING: Suspected of damaging fertility or the unborn child.

May cause damage to organs. **Read label before use.** Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Use personal protective equipment as required. Do not eat, drink or smoke when using this product. May cause drowsiness or dizziness. May cause respiratory irritation. Do not breathe vapours. Use only outdoors or in a well-ventilated area. Wash hands thoroughly after handling. **FIRST AID:** Call a POISON CENTER (0800 764 766) or doctor/physician if you feel unwell. If exposed or concerned get medical advice/attention. **IF INHALED:** Remove to fresh air and keep at rest in a position comfortable for breathing. **STORAGE:** Store locked up. Store in a well-ventilated place. Keep container tightly closed. **DISPOSAL:** Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility.

ACVM Registration No. A011436
See www.foodsafety.govt.nz for registration conditions

CATOLYST®

CATOLYST® controls coccidiosis caused by *Eimeria bovis* or *Eimeria zuernii* in young cattle.

Coccidiosis typically occurs in cattle less than a year old causing diarrhoea which is often blood stained. Recently weaned calves are particularly susceptible to outbreaks of clinical disease. CATOLYST® is effective in one dose because it attacks all stages of the parasite in the animal.

DIRECTIONS FOR USE By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label. Thoroughly shake container before use.

CATTLE DOSE: Give 3 mls of CATOLYST® per 10kg of bodyweight by mouth. For the treatment of clinical disease treat all affected and in contact animals. For prevention of coccidiosis on farms with a known history treat prior to the expected onset of clinical signs. For prevention of coccidiosis in recently weaned calves treat at weaning time when meal feeding ceases.

WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Cattle producing meat and offal for human consumption must not be sold for slaughter during or within 56 days of the last treatment.

ACVM Registration No. A011436
See www.foodsafety.govt.nz for registration conditions

PLCAT1B-01
STORE below 25°C
B.
EXP.



Registered to Alleva
Animal Health Ltd
15 Calman Place,
Birkenhead, Auckland

CATOLYST® is a registered trademark of Alleva Animal Health



PEEL HERE