

WARNING:
READ LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

CORPORAL[®]

DUAL COMBINATION ORAL DRENCH

FOR THE CONTROL AND TREATMENT OF
INTERNAL PARASITES IN SHEEP AND CATTLE

Active ingredients: 90.6 mg/ml Oxfendazole
and 150 mg/ml Levamisole HCL

LOW DOSE – COMBINATION DRENCH

NET CONTENTS: 5L

PLCOR5F-01





FOR ANIMAL TREATMENT ONLY
WARNING DANGEROUS TO THE
ENVIRONMENT

Corporal is a low volume drench that contains oxfendazole as a member of the benzimidazoles and levamisole HCL as a member of the imidazothiazole family. It is effective against sensitive strains of the following parasites: Mature and immature Barber's Pole Worm (*Haemonchus*), Small Brown Stomach Worm (*Teladorsagia & Ostertagia*), Stomach Hair Worm and Black Scour Worm (*Trichostrongylus*), Thin-Necked Intestinal Worm (*Nematodirus*), Small Intestinal Worm (*Cooperia*), Intestinal Threadworm (*Strongyloides*), Hookworm (*Bunostomum*), Large Bowel Worm (*Oesophagostomum*), Large-Mouthed Bowel Worm (*Chabertia*), Whipworm (*Trichuris*), Tapeworm (*Moniezia*), Lungworm (*Dictyocaulus*) in cattle and sheep, type II Ostertagiosis in cattle and as an aid in the control of Liver Fluke in sheep.

Resistance can develop to any anthelmintic. It is advisable that a resistance test be conducted regularly when using any parasite treatment. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance.

PRECAUTION: DO NOT EXCEED THE STATED DOSE VOLUME AND FREQUENCY. DOSES OF 3 OR MORE TIMES THE RECOMMENDED CAN CAUSE SYMPTOMS OF LEVAMISOLE TOXICITY, SO ESTIMATE LIVeweIGHTS CAREFULLY. DEHYDRATED ANIMALS MAY BE MORE SUSCEPTIBLE TO TOXICITY. FATAL INTERACTIONS MAY OCCUR BETWEEN LEVAMISOLE AND ORGANOPHOSPHATE DIPS.

DIRECTIONS FOR USE: Shake well before use. Corporal is ready to use through standard drenching equipment. Correct drenching technique should be used. A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group. Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. For drenching sheep, use a gun calibrated to dose in 0.5mL increments. Rinse gun after use.

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

DOSE GUIDE CATTLE: 1ml per 20kg bodyweight
(4.5mg/kg oxfendazole and 7.5mg/kg Levamisole HCL)

| Liveweight | Dose | 5L pack treats |
|------------|--------|----------------|
| 50kg | 2.5mL | 2000 |
| 100kg | 5mL | 1000 |
| 150kg | 7.5mL | 666 |
| 200kg | 10mL | 500 |
| 250kg | 12.5mL | 400 |
| 300kg | 15mL | 333 |
| 350kg | 17.5mL | 285 |
| 400kg | 20mL | 250 |
| 450kg | 22.5mL | 222 |
| 500kg | 25mL | 200 |
| 550kg | 27.5mL | 181 |
| 600kg | 30mL | 166 |
| 650kg | 32.5mL | 153 |

Cattle in excess of 650 kg to be dosed at 1mL/20 kg bodyweight.

DOSE GUIDE SHEEP: 1ml/20kg bodyweight
(4.5mg/kg oxfendazole and 7.5mg/kg Levamisole HCL)
Do not dose lambs under 20kg of bodyweight.

| Liveweight | Dose | 5L pack treats |
|------------|-------|----------------|
| 20kg | 1.0mL | 5000 |
| 30kg | 1.5mL | 3333 |
| 40kg | 2.0mL | 2500 |
| 50kg | 2.5mL | 2000 |
| 60kg | 3.0mL | 1666 |
| 70kg | 3.5mL | 1428 |
| 80kg | 4.0mL | 1250 |

Sheep in excess of 80kg should be dosed at 1ml/20kg bodyweight.

WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

MEAT: Animals producing meat and offal for human consumption must not be sold for slaughter either during treatment or within 10 days of last treatment.


MILK: Milk intended for sale for human consumption must be discarded during treatment and for not less than 35 days following the last treatment.

EPA Approval No. HSR100615
See www.epa.govt.nz for approval conditions
ACVM Registration No. A11042
See www.foodsafety.govt.nz for registration conditions

STORE below 25°C B.

EXP.


SAFETY PRECAUTIONS AND DIRECTIONS: May be harmful if swallowed. Wash hands and exposed skin before meals and after use. Repeated exposure may cause skin allergy. Avoid skin contact. Levamisole HCL may possibly cause damage to genetic material. Oxfendazole may affect development and/or reproduction. Oxfendazole (liver and alimentary system) and Levamisole HCL (blood and haemopoetic system) may cause organ damage. Handle with care.
FIRST AID: If swallowed do NOT induce vomiting. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs remove contaminated clothing and flush skin and hair with running water. **ENVIRONMENTAL PROTECTION:** Toxic to aquatic organisms. Avoid contamination of any water supply with product or empty container. Harmful to terrestrial vertebrates. Avoid release to the environment. **Emergency Response:** In case of spillage, wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash the area with water and absorb with further inert material. Dispose of waste safely. **DISPOSAL:** Preferably dispose of the product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Burn empty container in an appropriate incinerator, if circumstances such as wind direction permit. Otherwise crush or puncture and bury in a suitable landfill. Do NOT use container for any other purpose.



FREE RECYCLING



agrecovery.co.nz



Registered to
Alleva Animal Health Ltd
15 Calman Place,
Birkenhead, Auckland

PLCOR5B-01

WARNING:
READ LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

C  **RPORAL** [®]

DUAL COMBINATION ORAL DRENCH

**FOR THE CONTROL AND TREATMENT OF
INTERNAL PARASITES IN SHEEP AND CATTLE**

**Active ingredients: 90.6 mg/ml Oxfendazole
and 150 mg/ml Levamisole HCL**

LOW DOSE – COMBINATION DRENCH

NET CONTENT: 20L



alleva
ANIMAL HEALTH

CORPORAL[®]

DUAL COMBINATION ORAL DRENCH

FOR ANIMAL TREATMENT ONLY

WARNING DANGEROUS TO THE ENVIRONMENT

Corporal is a low volume drench that contains oxfendazole as a member of the benzimidazoles and levamisole HCL as a member of the imidazothiazole family. It is effective against sensitive strains of the following parasites: Mature and immature Barber's Pole Worm (*Haemonchus*), Small Brown Stomach Worm (*Teladorsagia & Ostertagia*), Stomach Hair Worm and Black Scour Worm (*Trichostrongylus*), Thin-Necked Intestinal Worm (*Nematodirus*), Small Intestinal Worm (*Cooperia*), Intestinal Threadworm (*Strongyloides*), Hookworm (*Bunostomum*), Large Bowel Worm (*Oesophagostomum*), Large-Mouthed Bowel Worm (*Chabertia*), Whipworm (*Trichuris*), Tapeworm (*Moniezia*), Lungworm (*Dictyocaulus*) in cattle and sheep, type II Ostertagiosis in cattle and as an aid in the control of Liver Fluke in sheep.

Resistance can develop to any anthelmintic. It is advisable that a resistance test be conducted regularly when using any parasite treatment. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance.

PRECAUTION: DO NOT EXCEED THE STATED DOSE VOLUME AND FREQUENCY. DOSES OF 3 OR MORE TIMES THE RECOMMENDED CAN CAUSE SYMPTOMS OF LEVAMISOLE TOXICITY, SO ESTIMATE LIVeweIGHTS CAREFULLY. DEHYDRATED ANIMALS MAY BE MORE SUSCEPTIBLE TO TOXICITY. FATAL INTERACTIONS MAY OCCUR BETWEEN LEVAMISOLE AND ORGANOPHOSPHATE DIPS.

DIRECTIONS FOR USE: Shake well before use. Corporal is ready to use through standard drenching equipment. Correct drenching technique should be used. A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group. Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. For drenching sheep, use a gun calibrated to dose in 0.5mL increments. Rinse gun after use.

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

MEAT: Animals producing meat and offal for human consumption must not be sold for slaughter either during treatment or within 10 days of last treatment.

MILK: Milk intended for sale for human consumption must be discarded during treatment and for not less than 35 days following the last treatment.

EPA Approval No. HSR100615

See www.epa.govt.nz for approval conditions

ACVM Registration No. A11042

See www.foodsafety.govt.nz for registration conditions

STORE below 25°C

B.
EXP.

DOSE GUIDE CATTLE: 1ml per 20kg bodyweight

(4.5mg/kg oxfendazole and 7.5mg/kg Levamisole HCL)

| Liveweight | Dose | 20L pack treats |
|------------|--------|-----------------|
| 50kg | 2.5mL | 8000 |
| 100kg | 5mL | 4000 |
| 150kg | 7.5mL | 2666 |
| 200kg | 10mL | 2000 |
| 250kg | 12.5mL | 1600 |
| 300kg | 15mL | 1333 |
| 350kg | 17.5mL | 1142 |
| 400kg | 20mL | 1000 |
| 450kg | 22.5mL | 888 |
| 500kg | 25mL | 800 |
| 550kg | 27.5mL | 727 |
| 600kg | 30mL | 666 |
| 650kg | 32.5mL | 615 |

Cattle in excess of 650 kg to be dosed at 1mL/20 kg bodyweight.

DOSE GUIDE SHEEP: 1ml/20kg bodyweight (4.5mg/kg oxfendazole and 7.5mg/kg Levamisole HCL) Do not dose lambs under 20kg of bodyweight.

| Liveweight | Dose | 20L pack treats |
|------------|-------|-----------------|
| 20kg | 1.0mL | 20000 |
| 30kg | 1.5mL | 13333 |
| 40kg | 2.0mL | 10000 |
| 50kg | 2.5mL | 8000 |
| 60kg | 3.0mL | 6666 |
| 70kg | 3.5mL | 5714 |
| 80kg | 4.0mL | 5000 |

Sheep in excess of 80kg should be dosed at 1ml/20kg bodyweight.

SAFETY PRECAUTIONS AND DIRECTIONS: May be harmful if swallowed.

Wash hands and exposed skin before meals and after use. Repeated exposure may cause skin allergy. Avoid skin contact. Levamisole HCL may possibly cause damage to genetic material. Oxfendazole may affect development and/or reproduction. Oxfendazole (liver and alimentary system) and Levamisole HCL (blood and haemopoietic system) may cause organ damage. Handle with care.

FIRST AID: If swallowed do NOT induce vomiting. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs remove contaminated clothing and flush skin and hair with running water.

ENVIRONMENTAL PROTECTION: Toxic to aquatic organisms. Avoid contamination of any water supply with product or empty container. Harmful to terrestrial vertebrates. Avoid release to the environment. **Emergency Response:** In case of spillage, wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash the area with water and absorb with further inert material. Dispose of waste safely. **DISPOSAL:** Preferably dispose of the product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Burn empty container in an appropriate incinerator, if circumstances such as wind direction permit. Otherwise crush or puncture and bury in a suitable landfill. Do NOT use container for any other purpose.



Registered to
Alleva Animal Health Ltd
15 Calman Place,
Birkenhead, Auckland