

WARNING:
READ LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

EPRIASURE™

POUR-ON FOR CATTLE & DEER

FOR THE TREATMENT AND CONTROL OF INTERNAL
AND EXTERNAL PARASITES OF CATTLE AND
INTERNAL PARASITES OF DEER

Active ingredient: 5 mg/ml EPRINOMECTIN

NET CONTENTS: 5L

PLEPSF-01



EPRISURE™
POUR-ON FOR CATTLE & DEER

FOR ANIMAL TREATMENT ONLY

EPRISURE™ Pour-On contains 5 mg/ml eprinomectin. It is a broad spectrum anthelmintic for the treatment and control of internal parasites and external parasites in cattle and internal parasites in deer. Eprinomectin is a member of the macrocyclic lactones. It is effective against internal and external parasites sensitive to this family.

Cattle Gastrointestinal Roundworms: Adult, Immature and Inhibited L4: *Ostertagia ostertagi* and *Cooperia* spp. Adult and Immature: *Haemonchus contortus*, *Trichostrongylus axei*, *T.colubriformis*, *Oesophagostomum radiatum*, *Nematodirus helvetianus*, *Bunostomum phlebotomum*. Adult: *O.lyrata* and *Trichuris* spp. Resistance to macrocyclic lactones in *Cooperia* spp is widespread in New Zealand.

Cattle Lungworm: Adult and immature *Dictyocaulus viviparus*.

Cattle Lice: *Linognathus vituli*, *Haematopinus eurysternus*, *Solenopotes capillatus*, *Bovicola (Damalinia) bovis*.

Cattle Mites: *Chorioptes bovis*, *Sarcoptes scabiei* var *bovis*

Persistent Activity in Cattle:

EPRISURE™ Pour-On for Cattle and Deer when given at the recommended dosage of 1mL/10kg liveweight prevents reinfection of cattle with the following internal parasites for the time period indicated.

Deer Gastrointestinal Roundworms:

Adult and immature: *Trichostrongylus* spp. Adult: *Ostertagia -type* spp, *O.lyrata*, *O.leptospicularis*, *Spiculopteria spiculoptera*, *Sp.asymmetrica*, *Apertagia quadrispiculata*, *Oesophagostomum* spp.

Deer Lungworm: Adult and immature: *Dictyocaulus viviparus*

Resistance can develop to any chemical. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted regularly when using any parasite treatment.

DIRECTIONS FOR USE.

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Shake well before use. Ready for use through standard pour-on equipment. Check accuracy of pour-on applicator regularly. A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group. Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. Apply along the backline from shoulders to tail.

DOSE: 1ml per 10kg bodyweight
(0.5 mg eprinomectin per kg bodyweight)

Liveweight	Dose	5L pack treats
76-100kg	10 mL	500
101-125kg	12.5 mL	400
126-150kg	15 mL	333
151-175kg	17.5 mL	285
176-200kg	20 mL	250
201-250kg	25 mL	200
251-300kg	30 mL	166
301-350kg	35 mL	142
351-400kg	40 mL	125
401-450kg	45 mL	111
451-500kg	50 mL	100
501-550kg	55 mL	90
551-600kg	60 mL	83
601-650kg	65 mL	76

Animals in excess of 650 kg to be dosed at 5 mL per 50 kg bodyweight.

WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

CATTLE: DEER: Producing meat or offal for human consumption
Meat: Nil must not be sold for slaughter either during treatment
Milk: Nil or within 7 days of last treatment.

EPA Approval No. HSR001954

See www.epa.govt.nz for approval conditions

ACVM Registration No. A11052

See www.foodsafety.govt.nz for registration conditions

STORE below 25°C - Protect from low temperatures and light.
Close container when not in use.

B.

EXP.

Parasite*	Persistent Activity
<i>Ostertagia</i> spp.	28 days
<i>Cooperia</i> spp.	21 days
<i>Nematodirus helvetianus</i>	14 days
<i>Oesophagostomum radiatum</i>	28 days
<i>Dictyocaulus viviparus</i>	28 days
<i>Haemonchus</i> sp.	14 days
<i>Trichostrongylus</i> spp.	14 days

*The following parasite species are included within each of the relevant genera; *Ostertagia ostertagi*, *O. lyrata*, *O. leptospicularis*, *Cooperia oncophora*, *C. punctata*, *C. surnabada*, *Trichostrongylus axei*, *T.colubriformis*.

Obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

WARNING: Harmful if swallowed. Causes mild skin irritation. Causes eye irritation. May cause an allergic skin reaction. Suspected of causing genetic defects. Suspected of damaging fertility or the unborn child. May cause harm to breast-fed children. Very toxic to aquatic life. Toxic to terrestrial vertebrates. Very toxic to terrestrial invertebrates.

DANGER: Causes damage to organs.

HANDLING PRECAUTIONS: Do not breath vapours. Avoid contact with skin or eyes. Avoid contact during pregnancy / while nursing. Wear protective gloves, protective clothing, eye protection and face protection. Do not eat, drink or smoke when using this product. Contaminated work clothing should not be allowed out of the workplace. Wash hands thoroughly after handling. Avoid release into the environment. Collect spillage. Store locked up.

FIRST AID: If medical advice is needed have product container or label at hand.

IF SWALLOWED: Immediately call a POISON CENTER (0800 764 766) or doctor. Rinse mouth. IF SKIN IRRITATION OCCURS: get medical advice. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists get medical attention. IF ON SKIN: wash with plenty of soap and water. If skin irritation or rash occurs get medical advice. IF EXPOSED OR CONCERNED OR IF YOU FEEL UNWELL: Call a POISON CENTRE or doctor and get medical advice.

DISPOSAL: Dispose of the product by use or at an approved facility. Dispose of empty container at an approved facility, or burn in an appropriate incinerator. Avoid contamination of any water supply with chemical or empty container.



FREE RECYCLING



agrecovery.co.nz



Registered to
Alleva Animal Health Ltd
15 Calman Place,
Birkenhead, Auckland

PLEP5B-01

WARNING:
READ LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

EPRITMSURE

POUR-ON FOR CATTLE & DEER

FOR THE TREATMENT AND CONTROL OF INTERNAL
AND EXTERNAL PARASITES OF CATTLE AND
INTERNAL PARASITES OF DEER

Active ingredient: 5 mg/ml EPRINOMECTIN

NET CONTENTS: 20L



EPRISURE™ POUR-ON FOR CATTLE & DEER

FOR ANIMAL TREATMENT ONLY

EPRISURE™ Pour-On contains 5 mg/ml eprinomectin. It is a broad spectrum anthelmintic for the treatment and control of internal parasites and external parasites in cattle and internal parasites in deer. Eprinomectin is a member of the macrocyclic lactones. It is effective against internal and external parasites sensitive to this family.

Cattle Gastrointestinal Roundworms: Adult, Immature and Inhibited L4: *Ostertagia ostertagi* and *Cooperia* spp. Adult and Immature: *Haemonchus contortus*, *Trichostrongylus axei*, *T.colubriformis*, *Oesophagostomum radiatum*, *Nematodirus helvetianus*, *Bunostomum phlebotomum*. Adult: *O.lyrata* and *Trichuris* spp. Resistance to macrocyclic lactones in *Cooperia* spp is widespread in New Zealand.

Cattle Lungworm: Adult and immature *Dictyocaulus viviparus*.

Cattle Lice: *Linognathus vituli*, *Haematopinus eurysternus*, *Solenopotes capillatus*, *Bovicola (Damalinea) bovis*.

Cattle Mites: *Chorioptes bovis*, *Sarcoptes scabiei* var *bovis*

Persistent Activity in Cattle:

EPRISURE™ Pour-On for

Cattle and Deer when given at the recommended dosage of 1mL/10kg liveweight prevents reinfection of cattle with the following internal parasites for the time period indicated.

Parasite*	Persistent Activity
<i>Ostertagia</i> spp.	28 days
<i>Cooperia</i> spp.	21 days
<i>Nematodirus helvetianus</i>	14 days
<i>Oesophagostomum radiatum</i>	28 days
<i>Dictyocaulus viviparus</i>	28 days
<i>Haemonchus</i> sp.	14 days
<i>Trichostrongylus</i> spp.	14 days

*The following parasite species are included within each of the relevant genera; *Ostertagia ostertagi*, *O. lyrata*, *O. leptospicularis*, *Cooperia oncophora*, *C. punctata*, *C. surnabada*, *Trichostrongylus axei*, *T.colubriformis*.

Deer Gastrointestinal

Roundworms:

Adult and immature:

Trichostrongylus spp. Adult: *Ostertagia -type* spp, *O.lyrata*, *O.leptospicularis*, *Spiculoptera spiculoptera*, *Sp.asymmetrica*, *Apteragia quadrispiculata*, *Oesophagostomum* spp.

Deer Lungworm: Adult and immature: *Dictyocaulus viviparus*

Resistance can develop to any chemical. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted regularly when using any parasite treatment.

DIRECTIONS FOR USE.

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label. Shake well before use. Ready for use through standard pour-on equipment. Check accuracy of pour-on applicator regularly. A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group. Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. Apply along the backline from shoulders to tail.

DOSE: 1ml per 10kg bodyweight
(0.5 mg eprinomectin per kg bodyweight)

Liveweight	Dose	20L pack treats
76-100kg	10 mL	2000
101-125kg	12.5 mL	1600
126-150kg	15 mL	1332
151-175kg	17.5 mL	1140
176-200kg	20 mL	1000
201-250kg	25 mL	800
251-300kg	30 mL	664
301-350kg	35 mL	568
351-400kg	40 mL	500
401-450kg	45 mL	444
451-500kg	50 mL	400
501-550kg	55 mL	360
551-600kg	60 mL	333
601-650kg	65 mL	304

Animals in excess of 650 kg to be dosed at 5 mL per 50 kg bodyweight.

Obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

WARNING: Harmful if swallowed. Causes mild skin irritation. Causes eye irritation. May cause an allergic skin reaction. Suspected of causing genetic defects. Suspected of damaging fertility or the unborn child. May cause harm to breast-fed children. Very toxic to aquatic life.

Toxic to terrestrial vertebrates. Very toxic to terrestrial invertebrates.

DANGER: Causes damage to organs.

HANDLING PRECAUTIONS: Do not breath vapours. Avoid contact with skin or eyes. Avoid contact during pregnancy / while nursing. Wear protective gloves, protective clothing, eye protection and face protection. Do not eat, drink or smoke when using this product. Contaminated work clothing should not be allowed out of the workplace. Wash hands thoroughly after handling. Avoid release into the environment. Collect spillage. Store locked up.

FIRST AID: If medical advice is needed have product container or label at hand.

IF SWALLOWED: Immediately call a POISON CENTER (0800 764 766) or doctor. Rinse mouth. IF SKIN IRRITATION OCCURS: get medical advice. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists get medical attention.

IF ON SKIN: wash with plenty of soap and water. If skin irritation or rash occurs get medical advice. IF EXPOSED OR CONCERNED OR IF YOU FEEL UNWELL: Call a POISON CENTRE or doctor and get medical advice.

DISPOSAL: Dispose of the product by use or at an approved facility.

Dispose of empty container at an approved facility, or burn in an approved incinerator. Avoid contamination of any water supply with chemical or empty container.

WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

CATTLE:

Meat: Nil

Milk: Nil

DEER: Producing meat or offal for human consumption

must not be sold for slaughter either during treatment or within 7 days of last treatment.

EPA Approval No. HSR001954

See www.epa.govt.nz for approval conditions

ACVM Registration No. A11052

See www.foodsafety.govt.nz for registration conditions

STORE below 25°C

Protect from low temperatures and light.

Close container when not in use.

B.

EXP.



FREE RECYCLING



agrecovery.co.nz



Registered to
Alleva Animal Health Ltd
15 Calman Place,
Birkenhead, Auckland

